

Notification Date: 01 March 2024

UPDATE to GYN and Molecular Testing at SBMF

The implementation of GYN and Molecular testing is now scheduled to begin in the third quarter of 2024. This adjustment ensures that we have the requisite sample volumes for efficient processing of these samples. Additionally, it provides the necessary time for the completion of the build and testing of our client interfaces.

All testing will be performed onsite at our laboratory located at 3355 Douglas Road in South Bend. Attached for your reference is the test list with corresponding CPT codes as well as the collection and container information guide.

In addition, we are researching the expansion of our services at SBMF including but not limited to Flow Cytometry with Molecular testing. By bringing these tests in-house, we can offer our clients the convenience of receiving all results from one lab.





If you have further questions or would like to learn more about this testing, please contact your Account Executive; Laura Lehman, 574-309-2591 or Meghan Berkheiser, 574-286-0287.

Pap Testing w/ CPT Codes
ThinPrep Pap Only 88174
ThinPrep Pap/HPV High-Risk Co-Test (ages 30-65) 88174, 87624
ThinPrep Pap w/ Reflex to HPV High-Risk if ASC-US 88174, 88141, 87624
ThinPrep Pap w/ Reflex to HPV High-Risk if ASC-US, CT/NG 88174, 88141, 87624, 87491, 87591
ThinPrep Pap w/ Reflex to HPV High-Risk if ASC-US, CT/NG/ATV 88174, 88141, 87624, 87491, 87591, 87661
ThinPrep Pap/HPV High-Risk Co-Test w/ Reflex to HPV Genotyping 16 18/45 88174, 87624, 87625
ThinPrep Pap/HPV High-Risk Co-Test w/ Reflex to HPV Genotyping 16 18/45, CT/NG 88174, 87624, 87625, 87491, 87591
ThinPrep Pap/HPV High-Risk Co-Test w/ Reflex to HPV Genotyping 16 18/45, CT/NG/ATV 88174, 87624, 87625, 87491, 87591, 87661
HPV Genotyping 16 18/45 Only 87625

Molecular Testing
Chlamydia trachomatis & Neisseria gonorrhoeae (CT/NG) 87491, 87591
Candida vaginosis & Trichomonas vaginalis (CV/ATV) 87481, 87661
Bacterial vaginosis (BV) 81513
Trichomonas vaginalis (ATV) 87661
Mycoplasma genitalium (M. gen) 87563

Molecular Panels
Vaginitis
Bacterial vaginosis (BV) 81513
Candida vaginosis & Trichomonas vaginalis (CV/ATV) 87481, 87661
Vaginitis+
Bacterial vaginosis (BV) 81513
Candida vaginosis & Trichomonas vaginalis (CV/ATV) 87481, 87661
Chlamydia trachomatis & Neisseria gonorrhoeae (CT/NG) 87491, 87591
Vaginitis+ with M. gen (2 swabs)
Bacterial vaginosis (BV) 81513
Candida vaginosis & Trichomonas vaginalis (CV/ATV) 87481, 87661
Chlamydia trachomatis & Neisseria gonorrhoeae (CT/NG) 87491, 87591
Mycoplasma genitalium (M. gen) 87563
Cervicitis/Urethritis
Chlamydia trachomatis & Neisseria gonorrhoeae (CT/NG) 87491, 87591
Trichomonas vaginalis (ATV) 87661
Mycoplasma genitalium (M. gen) 87563
Chlamydia trachomatis & Neisseria gonorrhoeae (CT/NG) 87491, 87591
Trichomonas vaginalis (ATV) 87661

Hologic® Collection Devices for Laboratory Testing

COLLECTION KIT	CONTAINER	TEST NAMES
<p>Aptima® Multitest Swab</p> <p>Store specimen at <u>room temperature</u> after collection.</p>		<p>Aptima® Combo 2 Assay Chlamydia trachomatis & Neisseria gonorrhoeae</p> <p>Aptima® Mycoplasma genitalium Assay</p> <p>Aptima® Trichomonas vaginalis Assay</p> <p>Aptima® Bacterial vaginosis Assay</p> <p>Aptima® Candida vaginosis & Trichomonas vaginalis Assay</p>
<p style="text-align: center;">Aptima® Urine Specimen</p> <p>Patient should not urinate for at least 1 hour prior to specimen collection. Send only the Aptima® Urine Specimen Tube. <u>No urine cups will be accepted.</u></p> <ul style="list-style-type: none"> • Direct patient to provide first-catch urine (~20-30 mL of initial urine stream) into urine collection cup. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity. Female patients should not cleanse labial area prior to providing specimen. • Remove cap and transfer 2 mL of urine into urine specimen transport tube using the pipette provided. • The correct volume of urine has been added when the fluid level is between black fill lines on urine specimen transport tube label. • Discard pipette. Tightly screw cap onto tube. This is now known as the "processed urine specimen." • <u>Refrigerate</u> specimen after collection. 		<p><u>Female Urine</u></p> <p>Aptima® Combo 2 Assay Chlamydia trachomatis & Neisseria gonorrhoeae</p> <p>Aptima® Mycoplasma genitalium Assay</p> <p>Aptima® Trichomonas vaginalis Assay</p> <p><u>Male Urine</u></p> <p>Aptima® Combo 2 Assay Chlamydia trachomatis & Neisseria gonorrhoeae</p> <p>Aptima® Mycoplasma genitalium Assay</p> <p>Aptima® Trichomonas vaginalis Assay</p>
<p style="text-align: center;">Aptima® Unisex Swab</p> <p>Patient should not urinate for at least 1 hour prior to specimen collection.</p> <p><u>Male Urethral Swab Specimens</u></p> <ul style="list-style-type: none"> • Discard white cleaning swab. It is not needed for male specimen collection. • Insert blue specimen collection swab 2-4 cm into urethra. Gently rotate swab clockwise for 2-3 seconds. Withdraw swab. If the soft tip is touched, laid down, or dropped, discard and replace with a new Aptima Unisex Swab Specimen Collection Kit. • Immediately place swab into transport tube so black score line is at top of tube. Align the score line with top edge of tube and carefully break shaft. Discard top portion of shaft. Tightly screw cap onto tube. <p><u>Female Endocervical Swab Specimens</u></p> <ul style="list-style-type: none"> • Use white cleaning swab to remove excess mucus from cervical os and surrounding mucosa. Discard. • Insert blue collection swab into endocervical canal. Gently rotate swab clockwise for 10-30 seconds. Withdraw swab; avoid any contact with vaginal mucosa. If the soft tip is touched, laid down, or dropped, discard and get a new Aptima Unisex Swab Specimen Collection Kit. • While holding swab in hand, unscrew tube cap. Do not spill tube contents. If spilled, discard and replace with a new Aptima Unisex Swab Specimen Kit. • Immediately place swab into transport tube so black score line is at top of tube. Align the score line with top edge of tube and carefully break shaft. Discard top portion of shaft. Tightly screw cap onto tube. • <u>Refrigerate</u> specimen after collection. 		<p>Aptima® Combo 2 Assay Chlamydia trachomatis & Neisseria gonorrhoeae</p> <p>Aptima® Mycoplasma genitalium Assay</p> <p>Aptima® Trichomonas vaginalis Assay</p>
<p>ThinPrep® Pap Test Vial</p> <p>Store specimen at <u>room temperature</u> after collection.</p>		<p>Aptima® Combo 2 Assay Chlamydia trachomatis & Neisseria gonorrhoeae</p> <p>Aptima® Trichomonas vaginalis Assay</p> <p>Aptima® HPV Assay</p> <p>Aptima® HPV 16 18/45 Genotype Assay</p>